



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the *C. nanus*, Kroyer, if it is not identical with it. It is 1.125 mm. long. The cephalothorax is about as wide as it is long 1.875 mm.; the first abdominal segment is obcordate 1.5 long and 1.25 wide; and the second long and narrow 1.375 long and 0.375 wide. The cephalic bothria 0.25 diameter.

APRIL 10.

Mr. USELMA C. SMITH, in the chair.

Nine persons present.

APRIL 17.

Mr. THOMAS MEEHAN, Vice-President, in the chair.

Twenty-nine persons present.

The following papers were presented for publication :—

“Distinctive Characters of *Odontaspis littoralis*.” By Joseph Leidy, M. D.

“Parasitic Crustacea.” By Joseph Leidy, M. D.

Note on Eleonorite from Sevier Co., Arkansas.—Prof. GEORGE A. KOENIG submitted his identification of Eleonorite. This species occurs in cavities of Dufrenite and likewise intimately interlaminated with it. It is of deep blood red color and gives a yellowish streak. The habitus is prismatic columnar, the prisms showing strong vitreous lustre and pleochroism—light yellow, in one direction, deep red brown in a direction at right angles. On the very scant material at the author's disposal, no planes are sufficiently developed to allow of an identification of the crystallographic characters. The specific gravity was found = 2.949. The crystals can be heated in a matras to red heat without decrepitation, without change of color, lustre and shape, while yielding a strong condensation of water. Reactions for iron and phosphoric acids. The mineral is slowly dissolved in nitric acid, rapidly in hydrochloric acid. The iron is entirely *ferrie*. The water is completely expelled at 250°C. Owing to the scantiness of the material, the analysis was made with only 54 mg. of the mineral.

This quantity yielded: water = 8 mg; $\text{Mg}^2 \text{P}^2 \text{O}^7$ = 26.3 mg. $\text{Fe}^2 \text{O}^3$ = 26.8 mg.; $\text{Al}^2 \text{O}^3$ = 2 mg. In percentage

$\text{H}^2 \text{O}$	= 14.81
$\text{P}^2 \text{O}^5$	= 30.93
$\text{Fe}^2 \text{O}^3$	= 49.60
$\text{Al}^2 \text{O}^3$	= 4.50
	<hr/> 99.84

This furnishes the ratio

$$\begin{aligned} \text{P}^2\text{O}^5 : (\text{Fe, Al})^2\text{O}^3 : \text{H}^2\text{O} &= 1 : 1.625 : 3.78 \\ &= 2 : 3.23 : 7.56 \\ &= 2 : 3 : 8 \end{aligned}$$

hence the formula $\text{Fe}^6\text{P}^4\text{O}^{19} + 8 \text{H}^2\text{O}$. This is the formula established by A. Streng (*Zeitschr. f. Kryst. and Min.* Vol. 7, p. 398). Groth deduced from it the theoretical formula $\text{Fe}^3 (\text{HO})^3 (\text{PO}^4)^2 \cdot 2\frac{1}{2} \text{H}^2\text{O}$. (*Tabell. Uebers. d. Mineral.* Braunschweig 1882.) In view of the fact that the whole of the water escapes at 250°C . without change of color or structure, Groth's formula, assuming three molecules of hydroxyl, can not well be sustained. The water must be considered wholly as water of crystallization.

Eleonorite has heretofore been known but from one locality, the "Eleonore" iron mine near Giessen, in Germany. Some specimens are in Mr. C. S. Bement's collection, with which the author compared the mineral from this new locality. It was first announced by A. Nies in 1880 (*Ber. d. Oberhess. Ges. fuer Natur. u. Heilk.* No. 19.) and subsequently studied by A. Streng, who determined the symmetry as monosymmetric and established the formula given above.

APRIL 24.

Mr. JOHN H. REDFIELD, in the chair.

Twenty-nine persons present.

Samuel H. Friend, M. D. was elected a member.

The following were ordered to be printed:—